

NOW FDA-APPROVED FOR CATS!

In feline diabetes

The **right balance** of proven performance and unparalleled support is only available from one insulin.

**Vetsulin® (porcine insulin zinc suspension).
15+ years of successful use globally plus a
comprehensive portfolio of support tools.**

Vetsulin is the best place to start when it comes to cats with newly diagnosed diabetes. You want a therapy proven to work in this challenging disease. With a long track record internationally,* Vetsulin now provides veterinarians in the US with a reliable new therapy option. In addition to proven performance, Vetsulin comes with a whole host of support materials and service simply unavailable with any other insulin product. Vetsulin. Striking the right balance in feline diabetes care.

www.vetsulin.com



DIABETES VETSUPPORT: 877.783.4840

vetsulin®
(porcine insulin zinc suspension)

See next page for side effects, warnings, and contraindications.

* Vetsulin is known as Caninsulin® in Canada, Europe, and Australia. Caninsulin and Vetsulin are registered trademarks of Intervet Inc. or an affiliate.


© 2008 Intervet Inc. All rights reserved. CA-VET-8696CH6/08

Intervet Inc. • PO Box 318, 29160 Intervet Lane
Millsboro, Delaware 19966-0318 • USA
Phone: 800 441 8272 • www.intervetusa.com

 **Intervet**
Schering-Plough Animal Health



Transitioning Feline Patients with Diabetes Mellitus to Vetsulin[®] (porcine insulin zinc suspension)



Feeding Plan:

- Feed *ad libitum* or two meals daily (provide food just prior to administering insulin)
- A diet suitable for a diabetic cat may encourage remission:
 - High protein, low carbohydrate (preferably canned)

General Recommendations:

- Ideally, the blood glucose values will range between 120 and 350 mg/dL for the majority of the curve in a well regulated diabetic cat.
- The dose will most likely need to be adjusted until adequate regulation is achieved and may require future adjustments based on changes in weight or medical history.
- Dose adjustments should be based on clinical signs and evaluation of a serial blood glucose curve. Increases in dose should be made in 1 IU per injection increments (BID).
- Allow at least 2–4 weeks between dose changes (unless evidence of hypoglycemia).
- Educate the client on the need for and importance of using U-40 syringes with Vetsulin.
- Discard open Vetsulin 30 days after first broaching the vial. 2.5 mL vials of Vetsulin are available, and will last the average cat 2–3 weeks.

NADA 141-236, Approved by FDA

vetsulin®

(porcine insulin zinc suspension)

PRODUCT INFORMATION

CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

vetsulin® is a sterile aqueous zinc suspension of purified porcine insulin.

| | | | |
|-------------------|------------------------------|---------|-------------------------------------|
| Each mL contains: | purified porcine insulin | 40 IU | (30% amorphous and 70% crystalline) |
| | Zinc (as chloride) | 0.08 mg | |
| | Sodium acetate trihydrate | 1.36 mg | |
| | Sodium chloride | 7.0 mg | |
| | Methylparaben (preservative) | 1.0 mg | |

pH is adjusted with hydrochloric acid and/or sodium hydroxide.

INDICATION

vetsulin® (porcine insulin zinc suspension) is indicated for the reduction of hyperglycemia and hyperglycemia-associated clinical signs in dogs and cats with diabetes mellitus.

DOSAGE AND ADMINISTRATION

USE OF A SYRINGE OTHER THAN A U-40 SYRINGE WILL RESULT IN INCORRECT DOSING.

FOR SUBCUTANEOUS INJECTION IN DOGS AND CATS ONLY

vetsulin® should be mixed by gentle rolling of the vial prior to withdrawing the dose from the vial. Using a U-40 insulin syringe, the injection should be administered subcutaneously, 2 to 5 cm (3/4 to 2 in) from the dorsal midline, varying from behind the scapulae to the mid-lumbar region and alternating sides.

Always provide the Owner Information Sheet with each prescription.

Dogs

The initial recommended vetsulin® dose is 0.5 IU insulin/kg body weight. Initially, this dose should be given once daily concurrently with, or right after a meal.

The veterinarian should re-evaluate the dog at appropriate intervals and adjust the dose based on clinical signs, urinalysis results, and glucose curve values until adequate glycemic control has been attained. In the US clinical study, glycemic control was considered adequate if an acceptable blood glucose curve was achieved (reduction in hyperglycemia and a nadir of 60-160 mg/dL), clinical signs of hyperglycemia (polyuria, polydipsia, and ketonuria) were improved, and hypoglycemia (blood glucose <50 mg/dL) was avoided. Twice daily therapy should be initiated if the duration of insulin action is determined to be inadequate. If twice daily treatment is initiated, the two doses should be 25% less than the once daily dose required to attain an acceptable nadir. For example, if a dog receiving 20 units of vetsulin® once daily has an acceptable nadir but inadequate duration of activity, the vetsulin® dose should be changed to 15 units twice daily.

Further adjustments in dosage may be necessary with changes in the dog's diet, body weight, or concomitant medication, or if the dog develops concurrent infection, inflammation, neoplasia, or an additional endocrine or other medical disorder.

Cats

The initial recommended dose in cats is 1 to 2 IU per injection. The injections should be given twice daily at approximately 12 hour intervals. For cats fed twice daily, the injections should be given concurrently with, or right after each meal. For cats fed *ad libitum*, no change in feeding schedule is needed. The veterinarian should re-evaluate the cat at appropriate intervals and adjust the dose based on clinical signs, urinalysis results, and glucose curve values until adequate glycemic control has been attained.

Further adjustments in dosage may be necessary with changes in the cat's diet, body weight, or concomitant medication, or if the cat develops concurrent infection, inflammation, neoplasia, or an additional endocrine or other medical disorder.

CONTRAINDICATIONS

Dogs and cats known to have a systemic allergy to pork or pork products should not be treated with vetsulin®. vetsulin® is contraindicated during periods of hypoglycemia.

WARNINGS

User Safety: For use in animals only. Keep out of the reach of children. Avoid contact with eyes. In case of contact, immediately flush eyes with copious amounts of water for 15 minutes. Accidental injection may cause clinical hypoglycemia. In case of accidental injection, seek medical attention immediately. Exposure to product may induce a local or systemic allergic reaction in sensitized individuals.

Animal Safety: Owners should be advised to observe for signs of hypoglycemia (see Owner Information Sheet). Use of this product, even at established doses, has been associated with hypoglycemia. An animal with signs of hypoglycemia should be treated immediately. Glucose should be given orally or intravenously as dictated by clinical signs. Insulin should be temporarily withheld and, subsequently, the dosage should be adjusted, if indicated.

Any change in insulin should be made cautiously and only under a veterinarian's supervision. Changes in insulin strength, manufacturer, type, species (animal, human) or method of manufacture (rDNA versus animal-source insulin) may result in the need for a change in dosage.

Appropriate diagnostic tests should be performed to rule out endocrinopathies in pets that are difficult to regulate (e.g., hyperadrenocorticism in dogs and hyperthyroidism in cats).

PRECAUTIONS

Animals presenting with severe ketoacidosis, anorexia, lethargy, and/or vomiting should be stabilized with short-acting insulin and appropriate supportive therapy until their condition is stabilized. As with all insulin products, careful patient monitoring for hypoglycemia and hyperglycemia are essential to attain and maintain adequate glycemic control and prevent associated complications. Overdosage can result in profound hypoglycemia and death. Progestogens, certain endocrinopathies, and glucocorticoids can have an antagonistic effect on insulin activity. Intact bitches should be ovariectomized. Progestogen and glucocorticoid use should be avoided.

Drug Interactions:

In the US clinical effectiveness studies, dogs and cats received various medications while being treated with vetsulin® including antimicrobials, antivirals, antifungals, antihistamines, analgesics, anesthetics/tranquilizers, diuretics, bronchodilators, corticosteroids (cats), NSAIDs, thyroid hormone supplementation, hyperthyroid medication (methimazole), internal and external parasiticides, anti-emetics, dermatological topical treatments and oral supplements, ophthalmic preparations containing antimicrobials and antiinflammatories, and various vaccines. No medication interactions were reported. This drug was not studied in dogs receiving corticosteroids.

Reproductive Safety: The safety and effectiveness of vetsulin® in breeding, pregnant, and lactating dogs and cats has not been evaluated.

Use in puppies and kittens: The safety and effectiveness of vetsulin® in puppies and kittens has not been evaluated.

ADVERSE REACTIONS

Dogs

In the field effectiveness and safety study, 66 dogs were treated with vetsulin®. Sixty-two dogs were included in the assessment of safety. Hypoglycemia (defined as blood glucose <50 mg/dL) with or without associated clinical signs occurred in 35.5% (22/62) of the dogs at various times during the study. Clinical signs of hypoglycemia were generally mild in nature (described as weakness, lethargy, stumbling, falling down, and/or depression). Disorientation and collapse were reported less frequently and occurred in 16.1% (10/62) of the dogs. Two dogs had a seizure and one dog died during the seizure. Although never confirmed, the presumptive diagnosis was hypoglycemia-induced seizures. In the rest of the dogs, hypoglycemia resolved with appropriate therapy and adjustments in insulin dosage.

Seven owners recorded the following observations about the injection site on the home monitoring forms: swollen, painful, sore, and a bleb under the skin.

The following clinical observations occurred in the field study following treatment with vetsulin® and may be directly attributed to the drug or may be secondary to the diabetic state or other underlying conditions in the dogs: hematuria, vomiting, diarrhea, pancreatitis, non-specific hepatopathy/pancreatitis, development of cataracts, and urinary tract infections.

Cats

In a field effectiveness and safety study, safety data was reported for 78 cats receiving vetsulin®. Hypoglycemia (defined as blood glucose <50 mg/dL) was reported in 61 cats (88 total incidences). Fifteen of the occurrences (involving 13 cats) were associated with clinical signs described as lethargy, diarrhea, decreased appetite/anorexia, vomiting, and hypothermia. One cat had seizures following accidental overdosing by the owner and again during the subsequent dose adjustment period. The cat responded to supportive therapy and had no further hypoglycemic episodes. In all cases of hypoglycemia, the clinical signs resolved following symptomatic treatment and/or dose adjustment.

Polyneuropathy was reported in 4 cats. Two injection site reactions were reported: one as a mildly thickened subcutaneous tissue reaction and the second as a mild bruising.

The following clinical observations occurred in the field study following treatment with vetsulin® and may be directly attributed to the drug or may be secondary to the diabetic state or other underlying conditions in the cats: vomiting, lethargy, diarrhea, decreased appetite/anorexia, pancreatitis, dermal events, respiratory disease, urinary tract disorder, renal disease, dehydration, weight loss, polydipsia, polyuria, behavioral change, and ocular discharge/conjunctivitis.

In a smaller field effectiveness and safety study, 14 cats were treated with vetsulin®. Hypoglycemia was reported in 6 cats (8 total occurrences). Lethargy not associated with hypoglycemia was reported in 4 cats (6 total occurrences). The following clinical observations occurred in the field study following treatment with vetsulin® and may be directly attributed to the drug or may be secondary to the diabetic state or other underlying conditions in the cats: foul odor to stool, diarrhea, dull coat, rapid, shallow breathing, stiff gate in rear, gallop rhythm, and pruritus with alopecia.

During the 1998-2007 period, the following adverse events in 50 cats treated with porcine insulin zinc suspension were reported to Intervet International and Intervet Inc.: Death, seizures, lack of effectiveness/dysregulation, hypoglycemia, allergic or skin reaction, lethargy, vomiting/diarrhea, injection pain, hyperthermia, nystagmus, PU/PD, and abnormal behavior.

To report adverse reactions, call 1-800-345-4735.

GENERAL PHARMACOLOGY

Porcine insulin is similar in amino acid structure to canine insulin. vetsulin® is classified as intermediate acting insulin. In dogs, vetsulin® has two peaks of activity following subcutaneous administration (the first at around 4 hours and the second at around 11 hours) (1). The duration of activity varies between 14 and 24 hours (1). In cats, the peak activity following subcutaneous administration of vetsulin® occurs between 1.5 and 8 hours (2), and the duration of activity varies between 8 and 12 hours (2). The peak(s), duration of activity and dose required to adequately control diabetic signs will vary between individuals.

EFFECTIVENESS

Dogs

A total of 66 client-owned dogs were enrolled in and 53 completed the effectiveness and safety field study. The dogs completing the study included 22 breeds of purebred and various mixed breed dogs ranging in age from 4.8 to 14 years, and ranging in weight from 4.2 to 51.3 kg. Of the dogs completing the study, 25 were spayed females and 28 were male (21 neutered and 7 intact).

Dogs were started on vetsulin® at a dose of 1 IU/kg plus a body weight-dependent dose supplement once daily. The initial treatment time to reach acceptable glycemic control (Dose determination period) ranged from 5 to 151 days. Dogs were evaluated for treatment effectiveness three times at 30-day intervals (Study Period). The blood glucose curve means and mean nadirs were compared pre- and post-treatment to assess effectiveness. The blood glucose curve mean was reduced from 370 mg/dL pre-treatment to 151 mg/dL, 185 mg/dL, and 184 mg/dL at the three treatment period evaluations. The blood glucose mean nadir was reduced from 315 mg/dL pre-treatment to 93 mg/dL, 120 mg/dL, and 119 mg/dL at the three treatment period evaluations. Sixty days after an adequate vetsulin® dose was initially established, 94%, 96% and 83% of study dogs experienced a reduction in polyuria, polydipsia, and ketonuria, respectively. Investigators reported adequate glycemic control an average of 81% of the time during the Study Period.

Cats

A total of 85 client-owned cats (53 males and 25 females—all neutered) of various breeds were enrolled in a 60 day field effectiveness and safety study with continued use up to Day 180. Seven cats were removed from the study prior to the Day 7 evaluation. The remaining cats ranged in age from 3 to 17.5 years and in weight from 1.9 to 10.8 kg. Seventy-two cats completed the study to Day 60 and 66 cats completed to Day 180. The cats were started on vetsulin® at an initial dose of 1 to 2 IU insulin twice daily. Scheduled evaluations occurred at Days 7, 14, 30, 60, and 180. Dose adjustments were allowed at and between the evaluations. Effectiveness was based on blood glucose curve mean, blood glucose nadir and improvement in clinical signs. Blood glucose curve means decreased from 394 mg/dL on Day 0 to 217 mg/dL on Day 60. The mean blood glucose nadir decreased from 343 mg/dL on Day 0 to 146 mg/dL on Day 60.

Fourteen client-owned cats (10 males and 4 females—all neutered) of various breeds were enrolled in a 60 day effectiveness and safety field study. The cats ranged in age from 5 to 14 years and in weight from 3.40 to 6.97 kg. Twelve cats completed the study. The cats were started on vetsulin® at an initial dose of 1 to 2 IU insulin twice daily. Scheduled evaluations occurred at Days 7, 14, 30, and 60. Dose adjustments were allowed at and between the evaluations. The blood glucose curve means decreased from 354 mg/dL on Day 0 to 162 mg/dL on Day 60. The mean blood glucose nadir decreased from 321 mg/dL on Day 0 to 99 mg/dL on Day 60.

HOW SUPPLIED

vetsulin® is supplied as a sterile injectable suspension in multidose vials containing either 2.5 mL or 10 mL of 40 IU/mL porcine insulin zinc suspension. Vials are supplied in cartons of one, 10 mL vial and cartons of ten, 2.5 mL vials.

STORAGE CONDITIONS

Store in an upright position under refrigeration at 2° to 8° C (36° to 46° F). Do not freeze. Protect from light.

Additional information about vetsulin® and diabetes mellitus can be found at www.vetsulin.com

Distributed by: INTERVET INC.
Millsboro, DE 19966

Made in Germany

01/08

References

- Graham P., Nash A., and McKellar Q. "Pharmacokinetics of porcine insulin zinc suspension in diabetic dogs" Journal of Small Animal Practice. 1997. Vol 38, October: 434-438.
- Martin G.J. and Rand J.S. "Pharmacokinetic and Pharmacodynamic Study of Caninsulin in Cats with Diabetes Mellitus" (2000), Internal Study Report.

vetsulin is a registered trademark of Intervet Inc. or an affiliate.
© 2008 Intervet Inc. All rights reserved. CA-VET-8649CH5/08

Intervet Inc.
PO Box 318, 29160 Intervet Lane
Millsboro, Delaware 19966-0318
USA
Phone: 800 441 8272
www.intervetusa.com